



Medical Dosages

**** Avoid an overdose. Make sure to check the concentration on your bottle. ****

Motrin Dosage (ibuprofen)

Do not administer to infants younger than 6 months old.

Administer every 6 hours as needed

<u>Pounds</u>	<u>Infant (50 mg./ 1.25 ml)</u>		<u>Children (100 mg / 5 ml)</u>	
6	0.7	ml	1.4	ml
7	0.8	ml	1.6	ml
8	0.9	ml	1.8	ml
9	1.0	ml	2.0	ml
10	1.1	ml	2.3	ml
11	1.3	ml	2.5	ml
12	1.4	ml	2.7	ml
13	1.5	ml	3.0	ml
14	1.6	ml	3.2	ml
15	1.7	ml	3.4	ml
16	1.8	ml	3.6	ml
17	1.9	ml	3.9	ml
18	2.0	ml	4.1	ml
19	2.2	ml	4.3	ml
20	2.3	ml	4.5	ml
25	2.8	ml	5.7	ml
30	3.4	ml	6.8	ml
35	4.0	ml	8.0	ml
40	4.5	ml	9.1	ml
45	5.1	ml	10.2	ml
50	5.7	ml	11.4	ml
55	6.3	ml	12.5	ml
60	6.8	ml	13.6	ml
65	7.4	ml	14.8	ml
70	8.0	ml	15.9	ml
75	8.5	ml	17.0	ml
80	9.1	ml	18.2	ml
85	9.7	ml	19.3	ml
90	10.0	ml	20.0	ml
95	10.0	ml	20.0	ml
100	10.0	ml	20.0	ml

Tylenol Dosage (acetaminophen)

Do not administer to infants younger than 2 months old.

Administer every 4 hours as needed

<u>Pounds</u>	<u>Infant (80 mg / 0.8 ml)</u>		<u>Children (160 mg / 5 ml)</u>	
6	0.4 ml		1.3 ml	
7	0.5 ml		1.5 ml	
8	0.5 ml		1.7 ml	
9	0.6 ml		1.9 ml	
10	0.7 ml		2.1 ml	
11	0.8 ml		2.3 ml	
12	0.8 ml		2.6 ml	
13	0.9 ml		2.8 ml	
14	1.0 ml		3.0 ml	
15	1.0 ml		3.2 ml	
16	1.1 ml		3.4 ml	
17	1.2 ml		3.6 ml	
18	1.2 ml		3.8 ml	
19	1.3 ml		4.0 ml	
20	1.4 ml		4.3 ml	
25	1.7 ml		5.3 ml	
30	2.0 ml		6.4 ml	
35	2.4 ml		7.5 ml	
40	2.7 ml		8.5 ml	
45	3.1 ml		9.6 ml	
50	3.4 ml		10.7 ml	
55	3.8 ml		11.7 ml	
60	4.1 ml		12.8 ml	
65	4.4 ml		13.8 ml	
70	4.8 ml		14.9 ml	
75	5.1 ml		16.0 ml	
80	5.1 ml		16.0 ml	
85	5.1 ml		16.0 ml	
90	5.1 ml		16.0 ml	
95	5.1 ml		16.0 ml	
100	5.1 ml		16.0 ml	